Appendix K. Landcover Classes

Water - All areas of open water or permanent ice/snow cover.

11. Open Water - areas of open water, generally with less than 25 percent or greater cover of water (per pixel).

Developed - areas characterized by high percentage (approximately 30% or greater) of constructed materials (e.g. asphalt, concrete, buildings, etc).

- 21. Low Intensity Residential Includes areas with a mixture of constructed materials and vegetation. Constructed materials account for 30-80 percent of the cover. Vegetation may account for 20 to 70 percent of the cover. These areas most commonly include single-family housing units. Population densities will be lower than in high intensity residential areas.
- 22. High Intensity Residential Includes heavily built up urban centers where people reside in high numbers. Examples include apartment complexes and row houses. Vegetation accounts for less than 20 percent of the cover. Constructed materials account for 80-100 percent of the cover.
- 23. Commercial/Industrial/Transportation Includes infrastructure (e.g. roads, railroads, etc.) and all highways and all developed areas not classified as High Intensity Residential.

Barren - Areas characterized by bare rock, gravel, sad, silt, clay, or other earthen material, with little or no "green" vegetation present regardless of its inherent ability to support life. Vegetation, if present, is more widely spaced and scrubby than that in the "green" vegetated categories; lichen cover may be extensive.

- 31. Bare Rock/Sand/Clay Perennially barren areas of bedrock, desert, pavement, scarps, talus, slides, volcanic material, glacial debris, and other accumulations of earthen material.
- 32. Quarries/Strip Mines/Gravel Pits Areas of extractive mining activities with significant surface expression.
- 33. Transitional Areas of sparse vegetative cover (less than 25 percent that are dynamically changing from one land cover to another, often because of land use activities. Examples include forest clearcuts, a transition phase between forest and agricultural land, the temporary clearing of vegetation, and changes due to natural causes (e.g. fire, flood, etc.)

Forested Upland - Areas characterized by tree cover (natural or semi-natural woody vegetation, generally greater than 6 meters tall); Tree canopy accounts for 25-100 percent of the cover.

- 41. Deciduous Forest Areas dominated by trees where 75 percent or more of the tree species shed foliage simultaneously in response to seasonal change.
- 42. Evergreen Forest Areas characterized by trees where 75 percent or more of the tree species maintain their leaves all year. Canopy is never without green foliage.
- 43. Mixed Forest Areas dominated by trees where neither deciduous nor evergreen species represent more than 75 percent of the cover present.

Planted/Cultivated - Areas characterized by herbaceous vegetation that has been planted or is intensively managed for the production of food, feed, or fiber; or is maintained in developed settings for specific purposes. Herbaceous vegetation accounts for 75-100 percent of the cover.

VIRGINIA'S COMPREHENSIVE WILDLIFE CONSERVATION STRATEGY Appendix K — Landcover Classes

- 81. Pasture/Hay Areas of grasses, legumes, or grass-legume mixtures planted for livestock grazing or the production of seed or hay crops.
- 82. Row Crops Areas used for the production of crops, such as corn, soybeans, vegetables, tobacco, and cotton.
- 85. Urban/Recreational Grasses Vegetation (primarily grasses) planted in developed settings for recreation, erosion control, or aesthetic purposes. Examples include parks, lawns, golf courses, airport grasses, and industrial site grasses.

Wetlands - Areas where the soil or substrate is periodically saturated with or covered with water as defined by Cowardin et al.

- 91. Woody Wetlands Areas where forest or shrubland vegetation accounts for 25-100 percent of the cover and the soil or substrate is periodically saturated with or covered with water.
- 92. Emergent Herbaceous Wetlands Areas where perennial herbaceous vegetation accounts for 75-100 percent of the cover and the soil or substrate is periodically saturated with or covered with water.